

Ampulla of Vater

**Protocol applies to all intra-ampullary, peri-ampullary,
and mixed intra- and peri-ampullary carcinomas.**

*Protocol revision date: January 2005
Based on AJCC/UICC TNM, 6th edition*

Procedures

- **Cytology** (No Accompanying Checklist)
- **Incisional Biopsy** (No Accompanying Checklist)
- **Ampullectomy**
- **Pancreaticoduodenectomy (Whipple Resection)**

Author

Carolyn C. Compton, MD, PhD

Department of Pathology, McGill University, Montreal, Quebec, Canada
For the Members of the Cancer Committee, College of American Pathologists

© 2005. College of American Pathologists. All rights reserved.

The College does not permit reproduction of any substantial portion of these protocols without its written authorization. The College hereby authorizes use of these protocols by physicians and other health care providers in reporting on surgical specimens, in teaching, and in carrying out medical research for nonprofit purposes. This authorization does not extend to reproduction or other use of any substantial portion of these protocols for commercial purposes without the written consent of the College.

The College of American Pathologists offers these protocols to assist pathologists in providing clinically useful and relevant information when reporting results of surgical specimen examinations of surgical specimens. The College regards the reporting elements in the "Surgical Pathology Cancer Case Summary (Checklist)" portion of the protocols as essential elements of the pathology report. However, the manner in which these elements are reported is at the discretion of each specific pathologist, taking into account clinician preferences, institutional policies, and individual practice.

The College developed these protocols as an educational tool to assist pathologists in the useful reporting of relevant information. It did not issue the protocols for use in litigation, reimbursement, or other contexts. Nevertheless, the College recognizes that the protocols might be used by hospitals, attorneys, payers, and others. Indeed, effective January 1, 2004, the Commission on Cancer of the American College of Surgeons mandated the use of the checklist elements of the protocols as part of its Cancer Program Standards for Approved Cancer Programs. Therefore, it becomes even more important for pathologists to familiarize themselves with the document. At the same time, the College cautions that use of the protocols other than for their intended educational purpose may involve additional considerations that are beyond the scope of this document.

Summary of Changes to Checklist(s)

Protocol revision date: January 2005

The following changes have been made to the data elements of the checklist(s) since the January 2004 protocol revision.

Ampullectomy Checklist**Microscopic**

Tumor Size: An optional data element was added to record two additional dimensions in addition to the greatest dimension, as shown below

Tumor Size

Greatest dimension: ____ cm

*Additional dimensions: ____ x ____ cm

____ Cannot be determined (see Comment)

Surgical Pathology Cancer Case Summary (Checklist)

*Protocol revision date: January 2005
Applies to invasive carcinomas only
Based on AJCC/UICC TNM, 6th edition*

AMPULLA OF VATER: Ampullectomy

Patient name:

Surgical pathology number:

Note: Check 1 response unless otherwise indicated.

MACROSCOPIC**Tumor Site**

- Intra-ampullary
 Peri-ampullary
 Junction of ampullary and duodenal mucosa
 Not specified

MICROSCOPIC**Histologic Type**

- Adenocarcinoma (not otherwise characterized)
 Papillary adenocarcinoma
 Adenocarcinoma, intestinal type
 Adenocarcinoma, gastric foveolar type
 Mucinous adenocarcinoma
 Clear cell adenocarcinoma
 Signet-ring cell carcinoma
 Adenosquamous carcinoma
 Other (specify): _____
 Carcinoma, type cannot be determined

Histologic Grade

- Not applicable (histologic type not usually graded)
 GX: Cannot be assessed
 G1: Well differentiated
 G2: Moderately differentiated
 G3: Poorly differentiated
 G4: Undifferentiated
 Other (specify): _____

4

* Data elements **with asterisks** are **not required** for accreditation purposes for the Commission on Cancer. These elements may be clinically important, but are not yet validated or regularly used in patient management. Alternatively, the necessary data may not be available to the pathologist at the time of pathologic assessment of this specimen.

Tumor Size

Greatest dimension: ___ cm

*Additional dimensions: ___ x ___ cm

___ Cannot be determined (see Comment)

Pathologic Staging (pTNM)Primary Tumor (pT)

___ pTis: Carcinoma in situ

___ pT1: Tumor limited to ampulla of Vater or sphincter of Oddi

___ pT2: Tumor invades duodenal wall

Margins (check all that apply)

___ Not applicable

___ Cannot be assessed

___ Uninvolved by invasive carcinoma

Specify distance of invasive carcinoma to closest margin: ___ mm

___ Uninvolved by carcinoma in situ/adenoma

___ Involved by carcinoma in situ/adenoma

___ Involved by invasive carcinoma

***Additional Pathologic Findings (check all that apply)**

* ___ None identified

* ___ Dysplasia/adenoma

* ___ Ampullitis

* ___ Adenomyosis

* ___ Other (specify): _____

***Comment(s)**

* Data elements **with asterisks** are **not required** for accreditation purposes for the Commission on Cancer. These elements may be clinically important, but are not yet validated or regularly used in patient management. Alternatively, the necessary data may not be available to the pathologist at the time of pathologic assessment of this specimen.

Surgical Pathology Cancer Case Summary (Checklist)

*Protocol revision date: January 2005
Applies to invasive carcinomas only
Based on AJCC/UICC TNM, 6th edition*

**AMPULLA OF VATER: Pancreaticoduodenectomy
(Whipple Resection)**

Patient name:

Surgical pathology number:

Note: Check 1 response unless otherwise indicated.

MACROSCOPIC**Specimen Type**

- Pancreaticoduodenectomy
 Pancreaticoduodenectomy (pylorus sparing)
 Other (specify): _____
 Not specified

Tumor Site (check all that apply)

- Intra-ampullary
 Peri-ampullary
 Junction of ampullary and duodenal mucosa
 Not specified

Tumor Size

Greatest dimension: ___ cm

*Additional dimensions: ___ x ___ cm

 Cannot be determined (see Comment)**MICROSCOPIC****Histologic Type**

- Adenocarcinoma (not otherwise characterized)
 Papillary adenocarcinoma
 Adenocarcinoma, intestinal type
 Adenocarcinoma, gastric foveolar type
 Mucinous adenocarcinoma
 Clear cell adenocarcinoma
 Signet-ring cell carcinoma
 Adenosquamous carcinoma
 Other (specify): _____
 Carcinoma, type cannot be determined

6

* Data elements **with asterisks** are **not required** for accreditation purposes for the Commission on Cancer. These elements may be clinically important, but are not yet validated or regularly used in patient management. Alternatively, the necessary data may not be available to the pathologist at the time of pathologic assessment of this specimen.

Histologic Grade

- Not applicable
 GX: Cannot be assessed
 G1: Well differentiated
 G2: Moderately differentiated
 G3: Poorly differentiated
 G4: Undifferentiated
 Other (specify): _____

Pathologic Staging (pTNM)Primary Tumor (pT)

- pTX: Cannot be assessed
 pT0: No evidence of primary tumor
 pTis: Carcinoma in situ
 pT1: Tumor limited to ampulla of Vater or sphincter of Oddi
 pT2: Tumor invades duodenal wall
 pT3: Tumor invades pancreas
 pT4: Tumor invades peripancreatic soft tissues or other adjacent organs or structures

Regional Lymph Nodes (pN)

- pNX: Cannot be assessed
 pN0: No regional lymph node metastasis
 pN1: Regional lymph node metastasis
 Specify: Number examined: ____
 Number involved: ____

Distant Metastasis (pM)

- pMX: Cannot be assessed
 pM1: Distant metastasis
 *Specify site(s), if known: _____

Margins (check all that apply)

- Cannot be assessed
 Margins uninvolved by invasive carcinoma
 Distance of invasive carcinoma from closest margin: ____ mm
 Specify margin (if possible): _____
 Carcinoma in situ absent at pancreatic duct margin
 Carcinoma in situ present at pancreatic duct margin
 Carcinoma in situ absent at common bile duct margin
 Carcinoma in situ present at common bile duct margin
 Margins involved by invasive carcinoma
 Specify margin(s) (if possible): _____
 Not applicable

* Data elements **with asterisks** are **not required** for accreditation purposes for the Commission on Cancer. These elements may be clinically important, but are not yet validated or regularly used in patient management. Alternatively, the necessary data may not be available to the pathologist at the time of pathologic assessment of this specimen.

***Venous/Lymphatic (Large/Small Vessel) Invasion (V/L)**

- * Absent
- * Present
- * Indeterminate

***Perineural Invasion**

- * Absent
- * Present

***Additional Pathologic Findings (check all that apply)**

- * None identified
- * Dysplasia/adenoma
- * Ampullitis
- * Adenomyosis
- * Chronic pancreatitis
- * Acute pancreatitis
- * Gastritis
- * Other (specify): _____

***Comment(s)**

8 * Data elements **with asterisks** are **not required** for accreditation purposes for the Commission on Cancer. These elements may be clinically important, but are not yet validated or regularly used in patient management. Alternatively, the necessary data may not be available to the pathologist at the time of pathologic assessment of this specimen.

Background Documentation

Protocol revision date: January 2005

I. Cytologic Material

A. Clinical Information

1. Patient identification
 - a. Name
 - b. Identification number
 - c. Age (birth date)
 - d. Sex
2. Responsible physician(s)
3. Date of procedure
4. Other clinical information
 - a. Relevant history (eg, pancreatitis, familial adenomatous polyposis syndrome)
 - b. Relevant findings (eg, endoscopic, endoscopic retrograde cholangiopancreatography [ERCP], and/or imaging studies)
 - c. Clinical diagnosis
 - d. Procedure (eg, brushing, washing, other)
 - e. Operative findings
 - f. Anatomic site(s) of specimen(s)

B. Macroscopic Examination

1. Specimen
 - a. Unfixed/fixed (specify fixative)
 - b. Number of slides received, if appropriate
 - c. Quantity and appearance of fluid specimen, if appropriate
 - d. Other (eg, cytologic preparation from tissue)
 - e. Results of intraprocedural consultation
2. Material submitted for microscopic evaluation
3. Special studies, specify (eg, cytochemistry, immunocytochemistry)

C. Microscopic Evaluation

1. Adequacy of specimen (if unsatisfactory for evaluation, specify reason)
2. Tumor, if present (Note **A**)
 - a. Histologic type, if possible (Note **B**)
 - b. Histologic grade, if possible (Note **C**)
 - c. Other features (eg, necrosis)
3. Additional pathologic findings, if present
4. Results/status of special studies (specify)
5. Comments
 - a. Correlation with intraprocedural consultation, as appropriate
 - b. Correlation with other specimens, as appropriate
 - c. Correlation with clinical information, as appropriate

II. Incisional Biopsy

A. Clinical Information

1. Patient identification
 - a. Name
 - b. Identification number
 - c. Age (birth date)
 - d. Sex
2. Responsible physician(s)
3. Date of procedure

4. Other clinical information
 - a. Relevant history (eg, pancreatitis, familial adenomatous polyposis syndrome)
 - b. Relevant findings (eg, endoscopic, ERCP, and/or imaging studies)
 - c. Clinical diagnosis
 - d. Procedure (eg, endoscopic biopsy, ERCP biopsy)
 - e. Operative findings
 - f. Anatomic site(s) of specimen(s)

B. Macroscopic Examination

1. Specimen
 - a. Unfixed/fixed (specify fixative)
 - b. Number of pieces
 - c. Largest dimension of each piece
 - d. Results of intraoperative consultation
2. Submit entire specimen and frozen section tissue fragment(s) (unless saved for special studies)
3. Special studies (specify) (eg, histochemistry, immunohistochemistry)

C. Microscopic Evaluation

1. Tumor (Note **A**)
 - a. Histologic type (Note **B**)
 - b. Histologic grade (Note **C**)
 - c. Invasion
2. Additional pathologic findings, if present
 - a. Duodenitis
 - b. Adenoma
 - c. Other
3. Results/status of special studies (specify)
4. Comments
 - a. Correlation with intraoperative consultation, as appropriate
 - b. Correlation with other specimens, as appropriate
 - c. Correlation with clinical information, as appropriate

III. Ampullectomy

A. Clinical Information

1. Patient identification
 - a. Name
 - b. Identification number
 - c. Age (birth date)
 - d. Sex
2. Responsible physician(s)
3. Date of procedure
4. Other clinical information
 - a. Relevant history (eg, pancreatitis, familial adenomatous polyposis syndrome)
 - b. Relevant findings (eg, endoscopic, ERCP, and/or imaging studies)
 - c. Clinical diagnosis
 - d. Procedure
 - e. Operative findings
 - f. Anatomic site(s) of specimen(s)

B. Macroscopic Examination

1. Specimen
 - a. Organ(s)/tissue(s) included (specify)
 - b. Unfixed/fixed (specify fixative)
 - c. Number of pieces
 - d. Dimensions (3)

- e. Orientation, if indicated by surgeon
- f. Results of intraoperative consultation
- 2. Tumor, if visible
 - a. Location
 - b. Configuration
 - c. Size (Note **D**)
 - d. Descriptive features (eg, color, consistency, necrosis, hemorrhage)
 - e. Distance from margin of resection
- 3. Tissues submitted for microscopic evaluation
 - a. Submit entire specimen, include margin
 - b. Frozen section tissue fragment(s) (unless saved for special studies)
- 4. Special studies (specify) (eg, histochemistry, immunohistochemistry, electron microscopy)

C. Microscopic Evaluation

- 1. Tumor (Note **A**)
 - a. Histologic type (Note **B**)
 - b. Histologic grade (Note **C**)
 - c. Extent of invasion (Note **D**)
 - d. Venous/lymphatic vessel invasion (Note **E**)
 - e. Perineural invasion (Note **F**)
- 2. Margins, as appropriate
- 3. Additional pathologic findings, if present
- 4. Results/status of special studies (specify)
- 5. Comments
 - a. Correlation with intraoperative consultation, as appropriate
 - b. Correlation with other specimens, as appropriate
 - c. Correlation with clinical information, as appropriate

IV. Whipple Resection

(Pancreaticoduodenectomy, Partial or Complete, With or Without Partial Gastrectomy)

A. Clinical Information

- 1. Patient identification
 - a. Name
 - b. Identification number
 - c. Age (birth date)
 - d. Sex
- 2. Responsible physician(s)
- 3. Date of procedure
- 4. Other clinical information
 - a. Relevant history (eg, pancreatitis, familial adenomatous polyposis syndrome)
 - b. Relevant findings (eg, endoscopic, ERCP, and/or imaging studies)
 - c. Clinical diagnosis
 - d. Procedure (eg, gastroduodenal pancreatectomy, partial or complete)
 - e. Operative findings
 - f. Anatomic site(s) of specimen(s)

B. Macroscopic Examination

- 1. Specimen
 - a. Organ(s)/tissue(s) included (specify)
 - b. Unfixed/fixed (specify fixative)
 - c. Number of pieces
 - d. Dimensions (measure attached tissues individually)

- e. Orientation, if indicated by surgeon
- f. Results of intraoperative consultation
- 2. Tumor
 - a. Location
 - b. Configuration
 - c. Size (Note **D**)
 - d. Descriptive features (eg, color, consistency, necrosis, hemorrhage)
 - e. Estimated extent of invasion (Note **D**)
- 3. Margins (Note **G**)
- 4. Regional lymph nodes (Note **D**)
- 5. Additional pathologic findings, if present
 - a. Common bile duct obstruction
 - b. Pancreatic duct obstruction
 - c. Pancreatitis
 - d. Other
- 6. Tissues submitted for microscopic evaluation
 - a. Carcinoma, including
 - (1) points of deepest penetration of surrounding structures
 - (2) points of deepest penetration of closest margins
 - (3) interface of tumor with adjacent tissues
 - b. Accessory papilla, if present
 - c. Margins
 - (1) distal pancreas
 - (2) common bile duct
 - (3) posterior pancreatic surface (deep radial margin)
 - (4) proximal (gastric or duodenal)
 - (5) distal (duodenal)
 - d. All lymph nodes (Note **D**)
 - (1) regional
 - (2) nonregional
 - e. Other lesions (eg, pseudocysts)
 - f. Pancreas uninvolved by tumor
 - g. Other tissue(s)/organ(s)
 - h. Frozen section tissue fragment(s) (unless saved for special studies)
- 7. Special studies (specify) (eg, histochemistry, immunohistochemistry, electron microscopy)

C. Microscopic Evaluation

- 1. Tumor (Note **A**)
 - a. Histologic type (Note **B**)
 - b. Histologic grade (Note **C**)
 - c. Extent of invasion (Note **D**)
 - d. Venous/lymphatic vessel invasion (Note **E**)
 - e. Perineural invasion (Note **F**)
- 2. Margins (Note **G**)
 - a. Distal pancreas
 - b. Common bile duct
 - c. Posterior pancreatic surface (deep radial margin)
 - d. Proximal (gastric)
 - e. Distal (duodenal)
- 3. Regional lymph nodes (Note **D**)
 - a. Number
 - b. Number with metastases
- 4. Distant metastasis (specify site)

5. Additional pathologic findings, if present
 - a. Chronic pancreatitis
 - b. Dysplasia
 - c. Metaplasia
 - d. *Helicobacter pylori* gastritis (Note H)
 - e. Other
6. Other tissue(s)/organ(s)
7. Results/status of special studies (specify)
8. Comments
 - a. Correlation with intraoperative consultation, as appropriate
 - b. Correlation with other specimens, as appropriate
 - c. Correlation with clinical information, as appropriate

Explanatory Notes

A. Application

The ampulla of Vater is a complex structure that represents the confluence of the distal common bile duct and main pancreatic duct, which traverses the duodenal wall and opens into the duodenal lumen through a small mucosal elevation, the duodenal papilla. Tumors of the ampulla of Vater include tumors arising in the ampulla (intra-ampullary type), tumors arising on the ampulla (peri-ampullary type), and tumors arising at the junction of the mucosa of the ampulla with that of the papilla or involving both the intra-ampullary and peri-ampullary region of the duodenum (mixed type). Thus, tumors of the ampulla of Vater may show biliary and/or intestinal features. Tumors may be exophytic or ulcerated. The origin of the tumor may be difficult, and occasionally impossible, to determine. This protocol applies to all primary carcinomas of the ampulla of Vater including those showing focal endocrine differentiation, but it does not apply to carcinoid tumors, gastrointestinal stromal tumors, or lymphomas.^{1,2}

B. Histologic Type

This protocol applies to the following histologic classification but does not preclude the use of other histologic types or systems of classification. A modified classification of carcinomas of the extrahepatic bile ducts published by the World Health Organization (WHO) that is applicable to the ampulla of Vater is as follows.³

WHO Classification of Ampullary Carcinoma[#]

Adenocarcinoma
 Papillary adenocarcinoma[#]
 Adenocarcinoma, intestinal type
 Adenocarcinoma, gastric foveolar type
 Mucinous adenocarcinoma
 Clear cell adenocarcinoma
 Signet-ring cell carcinoma^{##}
 Adenosquamous carcinoma
 Squamous cell carcinoma
 Small cell carcinoma^{###}
 Large cell neuroendocrine carcinoma
 Undifferentiated carcinoma^{###}
 Biliary cystadenocarcinoma

The term “carcinoma, NOS (not otherwise specified)” is not part of the WHO classification.

Ampullary tumors of the papillary histologic type have been shown to have a favorable prognosis compared to tumors of non-papillary histologic types.^{4,5}

Signet-ring cell carcinomas are, by convention, classified as poorly differentiated (grade 3/4) adenocarcinomas, and poor differentiation has been shown to be an adverse prognostic factor for ampullary carcinomas (see below).^{5,6}

Small cell carcinomas and undifferentiated (histologic type) carcinomas are assigned grade 4 (see below).

C. Histologic Grade

For non-papillary adenocarcinomas, the following grading system is suggested:

- GX Grade cannot be assessed
- G1 Well differentiated (greater than 95% of tumor composed of glands)
- G2 Moderately differentiated (50% to 95% of tumor composed of glands)
- G3 Poorly differentiated[#] (49% or less of tumor composed of glands)

Poor differentiation has been shown to be an adverse prognostic factor.^{5,6}

Grade 4 carcinomas include both undifferentiated carcinomas (histologic type) and small cell carcinoma (high-grade neuroendocrine carcinomas) in the WHO classification (see above).

D. TNM and Stage Groupings

The TNM staging system for tumors of the ampulla of Vater of the American Joint Committee on Cancer (AJCC) and the International Union Against Cancer (UICC) is recommended and shown below.^{1,7} The post-resection prognosis of a patient with ampullary carcinoma is primarily determined by the anatomic extent of disease as defined by the TNM classification and stage groupings.⁶⁻¹¹

By AJCC/UICC convention, the designation “T” refers to a primary tumor that has not been previously treated. The symbol “p” refers to the pathologic classification of the TNM, as opposed to the clinical classification, and is based on gross and microscopic examination. pT entails a resection of the primary tumor or biopsy adequate to evaluate the highest pT category, pN entails removal of nodes adequate to validate lymph node metastasis, and pM implies microscopic examination of distant lesions. Clinical classification (cTNM) is usually carried out by the referring physician before treatment during initial evaluation of the patient or when pathologic classification is not possible.

Pathologic staging is usually performed after surgical resection of the primary tumor. Pathologic staging depends on pathologic documentation of the anatomic extent of disease, whether or not the primary tumor has been completely removed. If a biopsied tumor is not resected for any reason (eg, when technically unfeasible) and if the highest T and N categories or the M1 category of the tumor can be confirmed microscopically, the criteria for pathologic classification and staging have been satisfied without total removal of the primary cancer.

Primary Tumor (T)[#]

- TX Primary tumor cannot be assessed
 T0 No evidence of primary tumor
 Tis Carcinoma in situ
 T1 Tumor limited to the ampulla of Vater or sphincter of Oddi^{##}
 T2 Tumor invades the duodenal wall
 T3 Tumor invades pancreas
 T4 Tumor invades peripancreatic soft tissues or other adjacent organs or structures

[#] Although tumor size is not included in the TNM staging system for tumors of the ampulla of Vater, it has been shown to have independent prognostic significance as follows^{1,2}:

Tumor Size	5-Year Survival Rate (% ± SE)
Less than 2.5 cm	65%
2.5 cm or more	20%

The difference is statistically significant (p=0.039).

^{##} Invasion of the muscle of the sphincter of Oddi has been shown to be an adverse prognostic factor.⁵

Regional Lymph Nodes (N)[#]

- NX Regional lymph nodes cannot be assessed
 N0 No regional lymph node metastasis
 N1 Regional lymph node metastasis

[#] Regional lymph node metastases have been shown to have independent significance as an adverse prognostic factor.^{4,6,8,12}

The regional nodes may be subdivided as follows.

- Superior Lymph nodes superior to head and body of pancreas
 Inferior Lymph nodes inferior to head and body of pancreas
 Anterior Anterior pancreaticoduodenal, pyloric, and proximal mesenteric lymph nodes
 Posterior Posterior pancreaticoduodenal, common bile duct or pericholedochal, and proximal mesenteric nodes

The following lymph nodes are also considered regional: hepatic artery nodes, infrapyloric nodes, subpyloric nodes, celiac nodes, superior mesenteric nodes, retroperitoneal nodes, and lateral aortic nodes. Tumor involvement of other nodal groups is considered distant metastasis. Anatomic division of regional lymph nodes is not necessary, but separately submitted lymph nodes should be reported as submitted.¹

Regional Lymph Nodes (pN0): Isolated Tumor Cells

Isolated tumor cells (ITCs) are single cells or small clusters of cells not more than 0.2 mm in greatest dimension. Lymph nodes or distant sites with ITCs found by either histologic examination, immunohistochemistry, or nonmorphologic techniques (eg, flow cytometry, DNA analysis, polymerase chain reaction [PCR] amplification of a specific tumor marker) should be classified as N0 or M0, respectively. Specific denotation of the assigned N category is suggested as follows for cases in which ITCs are the only evidence of possible metastatic disease.^{13,14}

- pN0 No regional lymph node metastasis histologically, no examination for isolated tumor cells (ITCs)
- pN0(i-) No regional lymph node metastasis histologically, negative morphologic (any morphologic technique, including hematoxylin-eosin and immunohistochemistry) findings for ITCs
- pN0(i+) No regional lymph node metastasis histologically, positive morphologic (any morphologic technique, including hematoxylin-eosin and immunohistochemistry) findings for ITCs
- pN0(mol-) No regional lymph node metastasis histologically, negative nonmorphologic (molecular) findings for ITCs
- pN0(mol+) No regional lymph node metastasis histologically, positive nonmorphologic (molecular) findings for ITCs

Distant Metastasis (M)

- MX Presence of distant metastasis cannot be assessed
- M0 No distant metastasis
- M1 Distant metastasis

Stage Groupings

Stage 0	Tis	N0	M0
Stage IA	T1	N0	M0
Stage IB	T2	N0	M0
Stage IIA	T3	N0	M0
Stage IIB	T1	N1	M0
	T2	N1	M0
	T3	N1	M0
Stage III	T4	Any N	M0
Stage IV	Any T	Any N	M1

TNM Descriptors

For identification of special cases of TNM or pTNM classifications, the “m” suffix and “y,” “r,” and “a” prefixes are used. Although they do not affect the stage grouping, they indicate cases needing separate analysis.

The “m” suffix indicates the presence of multiple primary tumors in a single site and is recorded in parentheses: pT(m)NM.

The “y” prefix indicates those cases in which classification is performed during or following initial multimodality therapy (ie, neoadjuvant chemotherapy, radiation therapy, or both chemotherapy and radiation therapy). The cTNM or pTNM category is identified by a “y” prefix. The ycTNM or ypTNM categorizes the extent of tumor actually present at the time of that examination. The “y” categorization is not an estimate of tumor prior to multimodality therapy (ie, before initiation of neoadjuvant therapy).

The “r” prefix indicates a recurrent tumor when staged after a documented disease-free interval, and is identified by the “r” prefix: rTNM.

The “a” prefix designates the stage determined at autopsy: aTNM.

Additional Descriptors

Residual Tumor (R)

Tumor remaining in a patient after therapy with curative intent (eg, surgical resection for cure) is categorized by a system known as R classification, shown below.

RX	Presence of residual tumor cannot be assessed
R0	No residual tumor
R1	Microscopic residual tumor
R2	Macroscopic residual tumor

For the surgeon, the R classification may be useful to indicate the known or assumed status of the completeness of a surgical excision. For the pathologist, the R classification is relevant to the status of the margins of a surgical resection specimen. That is, tumor involving the resection margin on pathologic examination may be assumed to correspond to residual tumor in the patient and may be classified as macroscopic or microscopic according to the findings at the specimen margin(s).

Vessel Invasion

By AJCC/UICC convention, vessel invasion (lymphatic or venous) does not affect the T category indicating local extent of tumor unless specifically included in the definition of a T category. In all other cases, lymphatic and venous invasion by tumor are coded separately as follows.

Lymphatic Vessel Invasion (L)

LX	Lymphatic vessel invasion cannot be assessed
L0	No lymphatic vessel invasion
L1	Lymphatic vessel invasion

Venous Invasion (V)

VX	Venous invasion cannot be assessed
V0	No venous invasion
V1	Microscopic venous invasion
V2	Macroscopic venous invasion

E Venous/Lymphatic Invasion

Lymph and small blood vessel invasion has been shown to be an adverse prognostic factor.^{5,10}

F. Perineural Invasion

Perineural infiltration by tumor has been shown to be an adverse prognostic factor.⁵

G. Margins

Local recurrence from invasive carcinoma in the region of the pancreatic head most often occurs at the posterior margin of the pancreatic head. Since this is a critical margin, the protocol recommends inking the posterior retroperitoneal surface of the pancreas and submitting sections through the tumor at its closest approach to this margin. Local recurrence from intraductal tumor, however, is most likely to occur at a ductal resection

margin (ie, the main pancreatic duct and/or the common bile duct margin). Thus, complete en face sections through the distal pancreatic margin (representing the distal margin of the main pancreatic duct) and the margin of the common bile duct should also be taken. Involvement of resection margins has been shown to be an adverse prognostic factor.^{6,15}

H. Other Evaluation

Sections of gastric antrum may be evaluated for gastritis (eg, *Helicobacter pylori* gastritis, chemical gastritis).

References

1. Greene FL, Page DL, Fleming ID, et al, eds. *AJCC Cancer Staging Manual*. 6th ed. New York: Springer; 2002.
2. Albores-Saavedra J, Henson DE, Klimstra D. *Tumors of the Gallbladder and Extrahepatic Bile Ducts. Atlas of Tumor Pathology*. 3rd Series. Washington, DC: Armed Forces Institute of Pathology; 1999.
3. Albores-Saavedra J, Scoazec JC, Wittekind C, Sripa B. Carcinoma of the gallbladder and extrahepatic bile ducts. In: Hamilton SR, Aaltonen LA, eds. *World Health Organization Classification of Tumours. Pathology and Genetics of Tumours of the Digestive System*. Lyon: IARC Press; 2000:203-217.
4. Makipour H, Cooperman A, Danzi JT, Farmer RG. Carcinoma of the ampulla of Vater: Review of 38 cases with emphasis on treatment and prognostic factors. *Ann Surg*. 1976;183:341-344.
5. Griffanti-Bartoli F, Arnone GB, Ceppa P, Ravera G, Carrabetta S, Civalleri D. Malignant tumors in the head of the pancreas and the periampullary region, diagnostic and prognostic aspects. *Anticancer Res*. 1994;14:657-666.
6. Willett CG, Warshaw AL, Convery K, Compton CC. Patterns of failure after pancreaticoduodenectomy for ampullary carcinoma. *Surg Gynecol Obstet*. 1993;176:33-38.
7. Sobin LH, Wittekind C. *UICC TNM Classification of Malignant Tumours*. 6th ed. New York: Wiley-Liss; 2002.
8. Bakkevoid KE, Kambestad B. Staging of carcinoma of the pancreas and ampulla of Vater: tumor (T), lymph node (N), and distant metastasis as prognostic factors. *Int J Pancreatol*. 1995;17:249-259.
9. Bakkevoid KE, Kambestad B. Long-term survival following radical and palliative treatment of patients with carcinoma of the pancreas and papilla of Vater: the prognostic factors influencing the long-term results: a prospective multicentre study. *Eur J Surg Oncol*. 1993;19:147-161.
10. Mori K, Ikei S, Yamane T, et al. Pathological factors influencing survival of carcinoma of the ampulla of Vater. *Eur J Surg Oncol*. 1990;16:183-188.
11. Yamaguchi K, Enjoji M. Carcinoma of the ampulla of Vater: a clinicopathologic study and pathologic staging of 109 cases of carcinoma and 5 cases of adenoma. *Cancer*. 1987;59:506-515.
12. Delcore R, Connor CS, Thomas JH, Friesen SR, Hermreck AS. Significance of tumor spread in adenocarcinoma of the ampulla of Vater. *Am J Surg*. 1989;158:593-597.
13. Wittekind C, Greene FL, Henson DE, Hutter RVP, Sobin LH, eds. *TNM Supplement. A Commentary on Uniform Use*. 3rd ed. New York: Wiley-Liss; 2003.
14. Singletary SE, Greene FL, Sobin LH. Classification of isolated tumor cells: clarification of the 6th edition of the American Joint Committee on Cancer Staging Manual. *Cancer*. 2003 Dec 15;90(12):2740-2741.
15. Allema JH, Reinders ME, van Gulik TM, et al. Prognostic factors for survival after pancreaticoduodenectomy for patients with carcinoma of the pancreatic head region. *Cancer*. 1995;75:2069-2076.

Bibliography

- Bastid C, Bernard JP, Sarles H, et al. Mucinous ductal ectasia of the pancreas: a premalignant disease and a cause of obstructive pancreatitis. *Pancreas*. 1991;6:15-22.
- Braasch JW, Camer SJ. Periampullary carcinoma. *Med Clin North Am*. 1975;59:309-314.
- Conley CR, Scheithauer BW, van Heerden JA, et al. Diffuse intraductal papillary adenocarcinoma of the pancreas. *Ann Surg*. 1987; 205:246-249.
- Kawarada Y, Yano T, Yamamoto T, et al. Intraductal mucin-producing tumors of the pancreas. *Am J Gastroenterol*. 1992; 87:634-638.
- Klemptner J, Ridder GJ, Bektas H, Pichlmayr R. Surgery for exocrine pancreatic cancer: who are the 5- and 10-year survivors? *Oncology*. 1995;52:353-359.
- Knox RA, Kingston RD. Carcinoma of the ampulla of Vater. *Br J Surg*. 1986;73:72-73.
- Longnecker DS, Terhune PG. The case for parallel classification of biliary tract and pancreatic neoplasms. *Mod Pathol*. 1996;9:828-837.
- Milchgrub S, Campuzano M, Casillas J, et al. Intraductal carcinoma of the pancreas. *Cancer*. 1992;69:651-656.
- Morohoshi T, Kanda M, Asanuma K, et al. Intraductal papillary neoplasms of the pancreas: a clinicopathologic study of six patients. *Cancer*. 1989;64:1329-1335.
- Obara T, Saitoh Y, Maguchi H, et al. Multicentric development of a pancreatic intraductal carcinoma through atypical papillary hyperplasia. *Hum Pathol*. 1992;23:82-85.
- Shirai Y, Tsukada K, Ohtani T, et al. Carcinoma of the ampulla of Vater: histopathologic analysis of tumor spread in Whipple pancreatoduodenectomy specimens. *World J Surg*. 1995;19:102-107.
- Tian F, Myles J, Howard JM. Mucinous pancreatic ductal ectasia of latent malignancy: an emerging clinicopathologic entity. *Surgery*. 1992;111:109-113.
- Wise L, Pizzimbono C, Dehner LP. Periampullary cancer: a clinicopathologic study of 62 patients. *Am J Surg*. 1976;131:141-148.
- Wittekind C, Compton CC, Greene FL, Sobin LH. Residual tumor classification revisited. *Cancer*. 2002;94:2511-2516.
- Yamada M, Kozuka S, Yamao K, et al. Mucin-producing tumor of the pancreas. *Cancer*. 1991; 68:159-168.
- Yoshida J, Ozaki H, Yamamoto J, et al. Adenocarcinoma and concomitant intraductal papillary adenoma in the pancreas. *Jpn J Clin Oncol*. 1991;21:453-456.